

أسئلة الامتحان لدبلوم التعليم العام للمدارس الخاصة (ثنائية اللغة) الدور الأول مع نموذج الإجابة



تم تحميل هذا الملف من موقع المناهج العمانية

موقع فايلاتي ← المناهج العمانية ← الصف الثاني عشر ← كيمياء ← الفصل الأول ← الامتحان النهائي ← الملف

تاريخ إضافة الملف على موقع المناهج: 02:33:49 2025-10-08

ملفات اكتب للمعلم اكتب للطالب | اختبارات الكترونية | اختبارات | حلول | عروض بوربوينت | أوراق عمل
منهج انجليزي | ملخصات وتقارير | مذكرات وبنوك | الامتحان النهائي | للمدرس

المزيد من مادة
كيمياء:

التواصل الاجتماعي بحسب الصف الثاني عشر



صفحة المناهج
العمانية على
فيسبوك

المزيد من الملفات بحسب الصف الثاني عشر والمادة كيمياء في الفصل الأول

أسئلة الامتحان لدبلوم التعليم العام للمدارس الخاصة (ثنائية اللغة) الدور الأول مع نموذج الإجابة

1

إجابات أسئلة موضوعات الوحدة الثانية (الكيمياء الكهربائية)

2

أسئلة تدريبية في الوحدة الأولى

3

الخلاصة في الكيمياء (قوانين المنهج)

4

جميع مصطلحات الوجدتين الأولى والثانية

5

امتحان دبلوم التعليم العام للمدارس الخاصة (ثنائية اللغة)

الفصل الدراسي الأول - الدور الأول

للعام الدراسي ١٤٤٧/١٤٤٦ هـ - ٢٠٢٤ / ٢٠٢٥ م

- زمن الإجابة: ثلاث ساعات.
- الإجابة في الورقة نفسها.

- تنبيه:** • المادة: الكيمياء.
• الأسئلة في (١٦) صفحة.

تعليمات مهمة:

- يجب الحضور إلى قاعة الامتحان قبل عشر دقائق على الأقل من بدء زمن الامتحان.
- يجب إحضار أصل ما يثبت الهوية وإبرازها للعاملين بالامتحانات.
- يجب الالتزام بالزي (الدشداشة البيضاء والمصر أو الكمة للذكور) والزي المدرسي للطالبات ، ويستثنى من ذلك الدارسون من غير العمانيين بشرط الالتزام بالذوق العام، ويمنع على جميع المتقدمات ارتداء النقاب داخل المركز وقاعات الامتحان.
- يحظر على الممتحنين اصطحاب الهواتف النقالة وأجهزة النداء الآلي وآلات التصوير والحواسيب الشخصية والساعات الرقمية الذكية والآلات الحاسبة ذات الصفة التخزينية والمجلات والصحف والكتب الدراسية والدفاتر والمذكرات والحقائب اليدوية والآلات الحادة أو الأسلحة أيّاً كان نوعها وأي شيء له علاقة بالامتحان.
- يجب على الممتحن الامتثال لإجراءات التفتيش داخل المركز طوال أيام الامتحان.

- يجب على الممتحن التأكد من استلام دفتر امتحانه، مغلفاً بخلاف بلاستيكي شفاف وغير ممزق، وهو مسؤول عنه حتى يسلمه لمراقبي اللجنة بعد الانتهاء من الإجابة.
 - يجب الالتزام بضوابط إدارة امتحانات دبلوم التعليم العام وما في مستواه وأية مخالفة لهذه الضوابط تعرضك للتدابير والإجراءات والعقوبات المنصوص عليها بالقرار الوزاري رقم ٥٨٨ / ٢٠١٥.
 - يقوم المتقدم بالإجابة عن أسئلة الامتحان المقالية بقلم الحبر (الأزرق أو الأسود).
 - يقوم المتقدم بالإجابة عن أسئلة الاختيار من متعدد بتظليل الشكل () وفق النموذج الآتي:
 س - عاصمة سلطنة عمان هي: ☐ القاهرة ☐ الدوحة ☐ مسقط ☒ أبوظبي
ملاحظة: يتم تظليل الشكل () باستخدام القلم الرصاص وعند الخطأ، امسح بعناية لإجراء التغيير.
 صحيح ☒ غير صحيح ☐ ☐ ☐ ☐ ☒

مُسَوَّدَة، لَا يَتَمَّ تَصْحِيحُهَا

Do not write in this space

Do not write in this space


Question 1: Multiple Choice Items**(14 marks)**

There are 14 multiple-choice items worth one mark each.

Shade in the bubble (☐) next to the **correct** answer for each of the following items.

- 1) Which of the following statements is correct about an acidic buffer solution?
- ☐ It resists changes in the pH in alkaline regions.
 - ☐ Its pH does not depend on the ratio of $[HA] / [A^-]$.
 - ☐ It is a mixture of a weak acid and its conjugate base.
 - ☐ An example of an acidic buffer solution is NH_3 with ammonium chloride.

- 2) Consider the following reaction and statements for the forward reaction:



- I. HCO_3^- is an acid and H_2CO_3 its conjugate base.
- II. HCO_3^- is a base and H_2CO_3 its conjugate acid.
- III. $H_2PO_4^-$ is an acid and HPO_4^{2-} its conjugate base.
- IV. $H_2PO_4^-$ is a base and HPO_4^{2-} its conjugate acid

Which statements are correct?

- | | |
|----------------------------------|---------------------------------|
| ☐ I and III | ☐ I and IV |
| ☐ II and III | ☐ II and IV |

- 3) What is the solubility product K_{sp} in $\text{mol}^3 \text{dm}^{-9}$ for a saturated solution of calcium hydroxide $Ca(OH)_2$ that contains $[OH^-] = 4.47 \times 10^{-2} \text{ mol dm}^{-3}$?

- | | |
|---|---|
| ☐ 1.23×10^{-3} | ☐ 2.05×10^{-3} |
| ☐ 4.46×10^{-5} | ☐ 8.97×10^{-5} |

- 4) Which of the following solutions will not affect the solubility of Iron(II) carbonate ($FeCO_3$)?

- | | |
|-------------------------------------|------------------------------------|
| ☐ $MgCO_3(aq)$ | ☐ $NaCl(aq)$ |
| ☐ $K_2CO_3(aq)$ | ☐ $FeCl_2(aq)$ |

Do not write in this space

Question 1 continued

- 5) Which of the following terms describes the combining of two half-cells?
- ☐ Electrodes
 ☐ Electrolytic cell
☐ Salt bridge
 ☐ Voltage cell
- 6) Which one of the following is the value of an electrode potential for a half cell containing $\text{Ag(s)}/\text{Ag}^+(\text{aq})$ at 298 K and a concentration of $0.050 \text{ mol dm}^{-3}$ of $\text{Ag}^+(\text{aq})$ ions?
- ☐ 0.27 V
 ☐ 0.72 V
☐ 0.80 V
 ☐ 0.88 V
- 7) Which of the following options is correct about an electrochemical cell consisting of a $\text{Mn}^{2+} / \text{Mn}$ half - cell and a $\text{Sn}^{2+} / \text{Sn}$ half – cell?





	Positive electrode	Cell voltage
☐	Sn	–1.04
☐	Sn	+1.04
☐	Mn	–1.32
☐	Mn	+1.32

- 8) Which of the following reactions is feasible?
- ☐ $\text{Cu(s)} + \text{Zn}^{2+}(\text{aq}) \longrightarrow \text{Cu}^{2+}(\text{aq}) + \text{Zn(s)}$
☐ $\text{Pb(s)} + \text{Sn}^{2+}(\text{aq}) \longrightarrow \text{Pb}^{2+}(\text{aq}) + \text{Sn(s)}$
☐ $\text{Sn(s)} + \text{Mg}^{2+}(\text{aq}) \longrightarrow \text{Sn}^{2+}(\text{aq}) + \text{Mg(s)}$
☐ $\text{Fe(s)} + \text{Cu}^{2+}(\text{aq}) \longrightarrow \text{Fe}^{2+}(\text{aq}) + \text{Cu(s)}$





Do not write in this space

Question 1 continued

13) Which product is formed when phenol reacts with excess bromine water ($\text{Br}_2(\text{aq})$)?



14) Which of the following is the correct order of increasing acidity?



Question 2: Extended responses**(56 marks)**

Write your answer for each of the following questions in the space provided.
Be sure to show all your work, including the correct units where applicable.

- 15) A selection of solutions for buffer mixtures and K_a for the acid are listed below. Use them to answer the following questions.

Buffer mixture		K_a for acid / mol dm ⁻³
0.0100 mol dm ⁻³ acid	0.0200 mol dm ⁻³ conjugate base	
HCOOH	HCOO ⁻	1.6×10^{-4}
NH ₄ ⁺	NH ₃	5.6×10^{-10}
HNO ₂	NO ₂ ⁻	4.7×10^{-4}
C ₆ H ₅ COOH	C ₆ H ₅ COO ⁻	6.3×10^{-5}
H ₂ SO ₃	HSO ₃ ⁻	1.5×10^{-2}
H ₂ PO ₄ ⁻	HPO ₄ ²⁻	6.3×10^{-8}

- a. Define the following terms:

- (i) Conjugate acid-base pair.

_____ [1]

- (ii) Buffer solution.

_____ [1]

Do not write in this space

Question 2 continued

- b. Which one of the above acids has the highest value of pK_a ?

[1]


- c. Justify by calculating the pH, that C_6H_5COOH , $C_6H_5COO^-$ mixture is an acidic buffer solution.

[2]

- d. Why can a mixture of methanoic acid $HCOOH$ and sodium methanoate $HCOONa$ act as a buffer solution whereas a solution of sodium methanoate $HCOONa$ alone will not act as a buffer solution.

[1]

- e. The carbon dioxide-hydrogencarbonate ion system is an important buffer in the human body.



State to which direction the position of the equilibrium will shift when the H^+ ion concentration increases. Explain your answer.

[2]

Do not write in this space

Question 2 continued

16) A beaker contains a saturated solution of lead (II) chloride and a solid lead (II) chloride both are in an equilibrium. (K_{sp} for $\text{PbCl}_2 = 1.6 \times 10^{-5} \text{ mol}^3 \text{ dm}^{-9}$).

- a. Write equilibrium expressions for the solubility products for (PbCl_2).

_____ [1]

- b. Calculate the solubility in mol dm^{-3} of (PbCl_2).

_____ [1]

- c. Calculate the solubility in mol dm^{-3} of (PbCl_2) when $0.0500 \text{ mol dm}^{-3}$ of sodium chloride (NaCl) is added.

_____ [1]

- d. Explain the effect of adding (NaCl) to the solubility of (PbCl_2).

_____ [1]

Question 2 continued

17) The following table shows values of $E^\ominus$ for half cells represented by virtual symbols.

Use them to answer the following questions:

Half cells	$E^\ominus / \text{V}$
$\text{X}^{2+}(\text{aq}) / \text{X}(\text{s})$	+ 0.35
$\text{Y}^{2+}(\text{aq}) / \text{Y}(\text{s})$	– 0.26
$\text{Z}^{2+}(\text{aq}) / \text{Z}(\text{s})$	+ 0.79
$\text{M}^{2+}(\text{aq}) / \text{M}(\text{s})$	– 0.80
$\text{L}^{2+}(\text{aq}) / \text{L}(\text{s})$	– 0.18

a. Define the term standard electrode potential.

_____[1]

b. State two conditions used to measure the standard electrode potential.

_____[2]

c. Which metal ion is the strongest oxidizing agent?

_____[1]

d. Which metal has the highest tendency to lose its electrons?

_____[1]

Do not write in this space



Do not write in this space

Question 2 continued

- e. What is the direction of electrons flow for the two half cells $Y^{2+}(aq) / Y(s)$ and $X^{2+}(aq) / X(s)$. Write the final reaction equation.

[2]

- 18) Study the following cells that show the electrolysis of some electrolytes under standard conditions, to answer the questions below.



- a. For cell A:

- (i) What substance is formed at the anode?

[1]

- (ii) Write a half-equation for the reaction occurring at the cathode.

[1]

Do not write in this space

Question 2 continued

b. For cell B:

(i) Write a half-equation for the reaction occurring at the cathode?

_____ [1]


(ii) If we use diluted aqueous solution of potassium chloride, what substance(s) are formed at the anode during the electrolysis?

_____ [1]

Explain your answer.

_____ [1]

19) Study the following electrolysis cell under standard conditions. Then answer the questions:



a. Write the equation that describe the relationship between the Faraday constant and the Avogadro constant.

_____ [1]

Do not write in this space

Question 2 continued

- b. What is the type of reaction that occurs at the anode?

_____ [1]

- c. In which electrode will the silver be formed?

_____ [1]

- d. Calculate the quantity of charge that passed during 75 minutes of electrolysis

_____ [1]


- e. Calculate the mass of the silver formed.

($F = 96500 \text{ C mol}^{-1}$)

_____ [3]

Question 2 continued

- 20) The Born–Haber cycle below can be used to determine the lattice energy of calcium oxide, CaO.



Do not write in this space

- a. Define the term lattice energy.

_____ [1]

- b. What are the two properties of ions that affect the numerical magnitude of lattice energy?

_____ [2]

Do not write in this space

- c. Identify the enthalpy changes labelled (A) and (B) in the cycle:

(A): _____ [1]

(B): _____ [1]


- d. Calculate the lattice energy for calcium oxide.

 _____ [1]

- e. Explain why the lattice energy of calcium oxide is greater than the lattice energy of lithium fluoride.

 _____ [1]

- 21) The energy cycle below shows the enthalpy changes when magnesium chloride is dissolved in water:



- a. Define the term standard enthalpy change of hydration.

 _____ [1]

Question 2 continued

Do not write in this space

b. From the energy cycle for magnesium chloride:

(i) Identify species (**A**) in the cycle, including its physical state.

_____ [1]

(ii) Identify the enthalpy change labelled by the number (**1**) in the cycle.

_____ [1]

(iii) Calculate the enthalpy of hydration for the chloride ion given that:

Enthalpy change	Value/ kJ mol ⁻¹
$\Delta H^\theta_{\text{latt}}[\text{MgCl}_2]$	-2592
$\Delta H^\theta_{\text{sol}}[\text{MgCl}_2]$	-55
$\Delta H^\theta_{\text{hyd}}[\text{Mg}^{2+}]$	-1920


_____ [1]

c. Explain why the enthalpy change of hydration of Mg^{2+} is more exothermic than the enthalpy change of hydration of Ba^{2+} .

_____ [1]

Do not write in this space

22) Study the reactions scheme below, to answer the following questions:



a. For reaction 1:

(i) State the reagent (A).

_____ [1]

(ii) Name the compound produced.

_____ [1]

b. For reaction 2:

(i) State the reagent (B).

_____ [1]

(ii) Name the compound produced.

_____ [1]

Do not write in this space

c. **For reaction 3:**

Bromobenzene is produced. Which is more reactive?

☐ Bromoethane☐ Bromobenzene

(Shade your answer)

Explain your answer:

_____ [2]

d. **For reaction 4:**

Draw the structural formula of organic compound (C).



[1]

23) Study the reactions sequence below, to answer the following questions:



a. State the reagents required for reaction 1.

_____ [2]

b. Name compound 1.

_____ [1]

c. **Draw** and **name** the organic compound 3 produced in reaction 3.

_____ [2]

d. Explain the acidity of compound 2.

_____ [1]

[End of the Examination]

Do not write in this space

Standard electrode potentials

Electrode reaction	$E^{\ominus}/V$	Electrode reaction	$E^{\ominus}/V$
$F_2(g) + 2e^- \rightleftharpoons 2F^-(aq)$	+ 2.87	$SO_4^{2-}(aq) + 4H^+(aq) + 2e^- \rightleftharpoons SO_2(g) + 2H_2O(l)$	+ 0.17
$S_2O_8^{2-}(aq) + 2e^- \rightleftharpoons 2SO_4^{2-}(aq)$	+ 2.01		
$H_2O_2(l) + 2H^+(aq) + 2e^- \rightleftharpoons 2H_2O(l)$	+ 1.77	$Cu^{2+}(aq) + e^- \rightleftharpoons Cu^+(aq)$	+ 0.15
$Pb^{4+}(aq) + 2e^- \rightleftharpoons Pb^{2+}(aq)$	+ 1.69	$Sn^{4+}(aq) + 2e^- \rightleftharpoons Sn^{2+}(aq)$	+ 0.15
$MnO_4^-(aq) + 8H^+(aq) + 5e^- \rightleftharpoons Mn^{2+}(aq) + 4H_2O(l)$	+ 1.52	$S_4O_6^{2-}(aq) + 2e^- \rightleftharpoons 2S_2O_3^{2-}(aq)$	+ 0.09
$PbO_2(s) + 4H^+(aq) + 2e^- \rightleftharpoons Pb^{2+}(aq) + 2H_2O(l)$	+ 1.47	$2H^+(aq) + 2e^- \rightleftharpoons H_2(g)$	0.00
$Cl_2(g) + 2e^- \rightleftharpoons 2Cl^-(aq)$	+ 1.36	$Fe^{3+}(aq) + 3e^- \rightleftharpoons Fe(s)$	- 0.04
$Cr_2O_7^{2-}(aq) + 14H^+(aq) + 6e^- \rightleftharpoons 2Cr^{3+}(aq) + 7H_2O(l)$	+ 1.33	$Pb^{2+}(aq) + 2e^- \rightleftharpoons Pb(s)$	- 0.13
$O_2(g) + 4H^+(aq) + 4e^- \rightleftharpoons 2H_2O(l)$	+ 1.23	$Sn^{2+}(aq) + 2e^- \rightleftharpoons Sn(s)$	- 0.14
$Br_2(aq) + 2e^- \rightleftharpoons 2Br^-(aq)$	+ 1.07	$Ni^{2+}(aq) + 2e^- \rightleftharpoons Ni(s)$	- 0.25
$VO_2^+(aq) + 2H^+(aq) + e^- \rightleftharpoons VO^{2+}(aq) + H_2O(l)$	+ 1.00	$V^{3+}(aq) + e^- \rightleftharpoons V^{2+}(aq)$	- 0.26
$VO_3^-(aq) + 4H^+(aq) + e^- \rightleftharpoons VO^{2+}(aq) + 2H_2O(l)$	+ 1.00	$Co^{2+}(aq) + 2e^- \rightleftharpoons Co(s)$	- 0.28
$ClO^-(aq) + H_2O(l) + 2e^- \rightleftharpoons Cl^-(aq) + 2OH^-(aq)$	+ 0.89	$Fe^{2+}(aq) + 2e^- \rightleftharpoons Fe(s)$	- 0.44
$NO_3^-(aq) + 10H^+(aq) + 8e^- \rightleftharpoons NH_4^+(aq) + 3H_2O(l)$	+ 0.87	$Cr^{3+}(aq) + 3e^- \rightleftharpoons Cr(s)$	- 0.74
$NO_3^-(aq) + 2H^+(aq) + e^- \rightleftharpoons NO_2(g) + H_2O(l)$	+ 0.81	$Zn^{2+}(aq) + 2e^- \rightleftharpoons Zn(s)$	- 0.76
$Ag^+(aq) + e^- \rightleftharpoons Ag(s)$	+ 0.80	$2H_2O(l) + 2e^- \rightleftharpoons H_2(g) + 2OH^-(aq)$	- 0.83
$Fe^{3+}(aq) + e^- \rightleftharpoons Fe^{2+}(aq)$	+ 0.77	$Cr^{2+}(aq) + 2e^- \rightleftharpoons Cr(s)$	- 0.91
$I_2(s) + 2e^- \rightleftharpoons 2I^-(aq)$	+ 0.54	$Mn^{2+}(aq) + 2e^- \rightleftharpoons Mn(s)$	- 1.18
$Cu^+(aq) + e^- \rightleftharpoons Cu(s)$	+ 0.52	$V^{2+}(aq) + 2e^- \rightleftharpoons V(s)$	- 1.20
$O_2(g) + 2H_2O(l) + 4e^- \rightleftharpoons 4OH^-(aq)$	+ 0.40	$Mg^{2+}(aq) + 2e^- \rightleftharpoons Mg(s)$	- 2.38
$Cu^{2+}(aq) + 2e^- \rightleftharpoons Cu(s)$	+ 0.34	$Na^+(aq) + e^- \rightleftharpoons Na(s)$	- 2.71
$VO^{2+}(aq) + 2H^+(aq) + e^- \rightleftharpoons V^{3+}(aq) + H_2O(l)$	+ 0.34	$Ca^{2+}(aq) + 2e^- \rightleftharpoons Ca(s)$	- 2.87
		$K^+(aq) + e^- \rightleftharpoons K(s)$	- 2.92

Do not write in this space

Periodic Table

1 H 1.01	Atomic number Mass number Element symbol 11 Na 22.99																2 He 4.00
3 Li 6.941	4 Be 9.012															9 F 19.00	10 Ne 20.18
11 Na 22.99	12 Mg 24.31															17 Cl 35.45	18 Ar 40.00
19 K 39.10	20 Ca 40.08	21 Sc 44.96	22 Ti 47.88	23 V 50.94	24 Cr 52.00	25 Mn 54.94	26 Fe 55.85	27 Co 58.93	28 Ni 58.69	29 Cu 63.55	30 Zn 65.38	31 Ga 69.72	32 Ge 72.59	33 As 74.92	34 Se 78.96	35 Br 79.90	36 Kr 83.80
37 Rb 85.47	38 Sr 87.62	39 Y 88.91	40 Zr 91.22	41 Nb 92.91	42 Mo 95.94	43 Tc (98)	44 Ru 101.1	45 Rh 102.9	46 Pd 106.4	47 Ag 107.9	48 Cd 112.4	49 In 114.8	50 Sn 118.7	51 Sb 121.8	52 Te 127.6	53 I 126.9	54 Xe 131.3
55 Cs 132.9	56 Ba 137.3	57 La* 138.9	72 Hf 178.5	73 Ta 180.9	74 W 183.9	75 Re 186.2	76 Os 190.2	77 Ir 192.2	78 Pt 195.1	79 Au 197.0	80 Hg 200.6	81 Tl 204.4	82 Pb 207.2	83 Bi 209.0	84 Po (209)	85 At (210)	86 Rn (222)
87 Fr (223)	88 Ra 226	89 Ac (227)															
Lanthanides			58 Ce 140.1	59 Pr 140.9	60 Nd 144.2	61 Pm (145)	62 Sm 150.4	63 Eu 152.0	64 Gd 157.3	65 Tb 158.9	66 Dy 162.5	67 Ho 164.9	68 Er 167.3	69 Tm 168.9	70 Yb 173.0	71 Lu 175.0	
Actinides			90 Th 232.0	91 Pa (231)	92 U 238	93 Np (237)	94 Pu (244)	95 Am (243)	96 Cm (247)	97 Bk (247)	98 Cf (251)	99 Es (252)	100 Fm (257)	101 Md (258)	102 No (259)	103 Lr (260)	

Do not write in this space

Do not write in this space

Do not write in this space

مُسَوِّدَة

Do not write in this space

مُسَوِّدَة

Do not write in this space

MARKING GUIDE

GENERAL EDUCATION DIPLOMA
BILINGUAL PRIVATE SCHOOLS
SEMESTER ONE - FIRST SESSION

CHEMISTRY

2024 / 2025



Exam Specification for Chemistry (Grade 12 - Bilingual) – Semester One (2024/2025)

Topic of the Units	Weighting %	Multiple Choice (20%)				Extended Response (80%)				Total Mark		
		No. of Questions	Marks	Cognitive Levels			No. of Questions	Marks	Cognitive Levels			
				Knowing (30%)	Applying (50%)	Reasoning (20%)			Knowing (30%)		Applying (50%)	Reasoning (20%)
Quantitative Equilibria	23%	14	4	1	2	1	10	12	4	6	2	16
Electrochemistry Lattice energy	33%		4	1	2	1		19	5	10	4	23
	21%		3	1	2	-		12	4	6	2	15
Arenes , Phenols Total	23%	3	1	1	1	56	13	4	6	3	16	
	100 %	14	4	7	3		17	28	11	70		





Distribution of cognitive domains and marks.

Item	Mark	Unit	Cognitive domain	Output
1	1	Quantitative Equilibria	Knowing	1.1.g
2	1	Quantitative Equilibria	Application	1.1.b
3	1	Quantitative Equilibria	Application	1.1.n
4	1	Quantitative Equilibria	Reasoning	1.1.o
5	1	Electrochemistry	Knowing	2.1. c
6	1	Electrochemistry	Application	2.1. m
7	1	Electrochemistry	Application	2.1. g 2.1.a .ii
8	1	Electrochemistry	Reasoning	2.1.h.ii
9	1	Lattice energy	Knowing	3.1.d, c
10	1	Lattice energy	Application	3.1.a
11	1	Lattice energy	Application	3.2.c
12	1	Arenes and Phenols	Knowing	4.1.b
13	1	Arenes and Phenols	Application	4.2.b.ii
14	1	Arenes and Phenols	Reasoning	4.2.d
15.a	2	Quantitative Equilibria	Knowing	1.1.b, f
15.b	1	Quantitative Equilibria	Application	1.1.e
15.c	2	Quantitative Equilibria	Application	1.1.j
15.d	1	Quantitative Equilibria	Reasoning	1.1.h
15.e	2	Quantitative Equilibria	Knowing	1.1.i
16.a	1	Quantitative Equilibria	Application	1.1.m
16.b	1	Quantitative Equilibria	Application	1.1.n
16.c	1	Quantitative Equilibria	Application	1.1. p
16.d	1	Quantitative Equilibria	Reasoning	1.1.o
17.a	1	Electrochemistry	Knowing	2.1.a.i
17.b	2	Electrochemistry	Knowing	2.1.e
17.c	1	Electrochemistry	Application	2.1.j
17.d	1	Electrochemistry	Application	2.1.j
17.e	2	Electrochemistry	Reasoning	2.1.h.i
18.a.i	1	Electrochemistry	Applying	2.2.a
18.a.ii	1	Electrochemistry	Application	2.2.a
18.b.i	1	Electrochemistry	Application	2.2.a
18.b.ii	2	Electrochemistry	Reasoning	2.2.a
19.a	1	Electrochemistry	Knowing	2.2.b
19.b	1	Electrochemistry	Knowing	2.2.a
19.c	1	Electrochemistry	Application	2.1.c.v
19.d	1	Electrochemistry	Application	2.2.c.i
19.e	3	Electrochemistry	Application	2.2.c.ii
20.a	1	Lattice energy	Knowing	3.1.a

General Education Diploma, Semester One, First Session
Bilingual Private Schools, Chemistry, 2024/2025

20.b	2	Lattice energy	Knowing	3.1.g
20.c	2	Lattice energy	Application	3.1.e
20.d	1	Lattice energy	Application	3.1.f
20.e	1	Lattice energy	Reasoning	3.1.g
21.a	1	Lattice energy	Knowing	3.2.a
21.b	3	Lattice energy	Application	3.2.b, c
21.c	1	Lattice energy	Reasoning	3.2.d
22.a.i	1	Arenes and Phenols	Knowing	4.1.d
22.a.ii	1	Arenes and Phenols	Application	4.1.d
22.b.i	1	Arenes and Phenols	Knowing	4.1.c.i
22.b.ii	1	Arenes and Phenols	Application	4.1.c.i
22.c	2	Arenes and Phenols	Reasoning	4.1.e
22.d	1	Arenes and Phenols	Application	4.1.c.iv
23.a	2	Arenes and Phenols	Knowing	4.2.a.i
23.b	1	Arenes and Phenols	Application	4.1.b
23.c	2	Arenes and Phenols	Application	4.2.b.i
23.d	1	Arenes and Phenols	Reasoning	4.2.c




TOTAL PAGES: 6

MARKS: 70

Question One (14 Marks)

There are 14 multiple-choice items. Each correct answer is worth ONE mark.

Item	Correct option	
1	C	It is a mixture of a weak acid and its conjugate base.
2	C	II and III
3	C	$4.46 \times 10^{-5} \text{ mol}^3 \text{ dm}^{-9}$
4	B	NaCl(aq)
5	D	Voltage cell
6	B	0.72 V
7	B	$\text{Sn} \quad +1.04$
8	D	$\text{Fe(s)} + \text{Cu}^{2+}(\text{aq}) \rightarrow \text{Fe}^{2+}(\text{aq}) + \text{Cu(s)}$
9	C	It will be more exothermic going from chlorine to bromine to iodine.
10	A	$\text{K(s)} \rightarrow \text{K(g)}$
11	A	-73
12	D	The carbon atoms in benzene are sp^3 hybridized.
13	D	
14	A	$\text{CH}_3\text{CH}_2\text{OH} < \text{H}_2\text{O} < \text{C}_6\text{H}_5\text{OH}$





Question Two (56 Marks)

Part	Section	The answer	The mark
15	a. (i)	An acid-base pair on each side of an acid–base equilibrium equation that are related to each other by the difference of a hydrogen ion (proton).	1
	a. (ii)	A solution that minimizes changes in the pH when moderate amounts of acid or base are added.	1
	b	NH_4^+	1
	c	$K_a = \frac{[\text{H}^+][\text{C}_6\text{H}_5\text{COO}^-]}{[\text{C}_6\text{H}_5\text{COOH}]}, 6.3 \times 10^{-5} = \frac{[\text{H}^+](0.0200)}{(0.0100)}$ $[\text{H}^+] = 3.15 \times 10^{-5} \text{ mol dm}^{-3} \quad [1 \text{ mark}]$ $\text{pH} = -\log_{10} [\text{H}^+] = -\log_{10} [3.15 \times 10^{-5}] = 4.5 \quad [1 \text{ mark}]$ Or $\text{pH} = \text{p}K_a + \log \frac{(0.0200)}{(0.0100)} \quad [1 \text{ mark}]$ $\text{pH} = 4.5 \quad [1 \text{ mark}]$	2
	d	Because in the mixture of HCOOH and HCOONa the <u>large reserve supply</u> of HCOOH and HCOO^- ensures that the concentration of HCOOH and HCOO^- molecules in solution does not change significantly to remove the added H^+ or OH^- , so the pH changes very little. Or Because there is no a large reserve supply of HCOOH and HCOO^- in a solution of sodium methanoate HCOONa alone.	1
	e	When the H^+ ion concentration increases: • the position of <u>the equilibrium shifts to the left</u> [1 mark] • <u>H^+ ions combine with HCO_3^- ions</u> to form carbon dioxide and water until equilibrium is restored, keeping the pH constant. [1 mark]	2
16	a	$K_{sp} = [\text{Pb}^{2+}][\text{Cl}^-]^2$	1
	b	Let solubility be “s” $[\text{Pb}^{2+}] = s, [\text{Cl}^-] = 2s, K_{sp} = 4s^3$ $s = 1.59 \times 10^{-2} \text{ mol dm}^{-3}$ (1mark for the correct value)	1
	c	Ignore $[\text{Cl}^-]$ in PbCl_2 $K_{sp} = [\text{Pb}^{2+}][\text{Cl}^-]^2$ $1.6 \times 10^{-5} = [\text{Pb}^{2+}][0.0500]^2$ $[\text{Pb}^{2+}] = 6.4 \times 10^{-3} \text{ mol dm}^{-3}$ (1mark for the correct value)	1
	d	The chloride ion is common to both sodium chloride and lead(II) chloride. The addition of the common ion, Cl^- , will reduce the solubility of the lead(II) chloride because the added chloride ions shift the position of equilibrium to the left, and the lead(II) chloride is precipitated.	1

Part	Section	The answer	The mark
17	a	The voltage produced when a standard half – cell is connected to standard hydrogen electrode under standard conditions. Or: The potential of a cell composed of two electrodes as measured under standard conditions.	1
	b	1- Ion concentration 1.00 mol dm^{-3} 2- Temperature: 298 K Or 25 C° 3 - The pressure 1 atm or 101 k pa or 1.0 bar (any two correct answers get the marks)	2
	c	Z^{2+}	1
	d	M	1
	e	Electrons flow from Y^{2+} / Y half-cell to X^{2+} / X half cell Or from more negative value of E° to more positive value. Or Y metal will lose electrons to X^{2+} / X half-cell. Or X^{2+} ions will accept electrons from Y^{2+} / Y half-cell. Or The more positive pole attracts the negative electrons. Or Move from the strong reducing agent Y(s) to strong oxidizing agent X^{2+} . [1 mark] $X^{2+}(\text{aq}) + Y(\text{s}) \rightleftharpoons X(\text{s}) + Y^{2+}(\text{aq})$ [1 mark]	2
18	a.i	I_2 (Iodine)	1
	a.ii	$Zn^{2+}(\text{aq}) + 2e^{-} \rightarrow Zn(\text{s})$ (The full equation must be correct)	1
	b.i	$2H^{+}(\text{aq}) + 2e^{-} \rightarrow H_2(\text{g})$ Or $H^{+}(\text{aq}) + e^{-} \rightarrow 1/2H_2(\text{g})$ (The full equation must be correct)	1
	b.ii	mixture of oxygen and chlorine gases. Or oxygen gas. [1 mark] Because the relatively lower concentration of Cl^{-} ions allows OH^{-} ions to fall below Cl^{-} ions in the discharge series. Or Because the discharge potential of OH^{-} is less than Cl^{-} in the electrochemical series (discharge series). [1 mark]	2



Part	Section	The answer	The mark
19	a	$F = L.e$	1
	b	Oxidation	1
	c	The cathode	1
	d	$Q = It$ $= 0.30 \times 75 \times 60$ $= 1350 \text{ C}$ [1 mark]	1
	e	$\text{Ag}^+(\text{aq}) + \text{e}^- \rightleftharpoons \text{Ag}(\text{s})$ The number of coulombs required to deposit 1mole of product at the electrode : $= F1 \quad [1 \text{ mark}]$ $m = \frac{1350 \times 108}{1 \times 96500} \quad [1 \text{ mark}]$ $= 1.51 \text{ g} \quad [1 \text{ mark}]$	3


Part	Section	The answer	The mark
20	a	The energy change when 1 mole of an ionic compound is formed from its gaseous ions under standard conditions.	1
	b	- Ionic charge. [1 mark] - Ionic radius. [1 mark]	2
	c	(A): $\Delta H_f^\ominus$ Or enthalpy change of formation [1 mark] (B): $\text{EA}_2(\text{O})$ Or Second electron affinity [1 mark]	2
	d	$\Delta H_{latt}^\ominus = \Delta H_f^\ominus[\text{CaO}] - (\Delta H_{at}^\ominus[\text{Ca}] + \text{IE}_1[\text{Ca}] + \text{IE}_2[\text{Ca}] + \Delta H_{at}^\ominus[\text{O}] + \text{EA}_1[\text{O}] + \text{EA}_2[\text{O}])$ $= -3454 \text{ kJ mol}^{-1} \quad [1 \text{ mark}]$	1
	e	Calcium oxide has a greater lattice energy than lithium fluoride. The doubly charged Ca^{2+} and O^{2-} ions in calcium oxide attract each other more strongly than the singly charged Li^+ and F^- ions in lithium fluoride.	1



Part	Section	The answer	The mark
21	a	The enthalpy change when 1 mole of a specified gaseous ion dissolves in sufficient water to form a very dilute solution under standard conditions.	1
	b (i)	$\text{MgCl}_2(\text{aq})$	1
	b (ii)	$\Delta H_{\text{latt}}^{\ominus}[\text{MgCl}_2]$ Or lattice enthalpy	1
	b (iii)	$2\Delta H_{\text{hyd}}^{\ominus}[\text{Cl}^-] = \Delta H_{\text{latt}}^{\ominus}[\text{MgCl}_2] + \Delta H_{\text{sol}}^{\ominus}[\text{MgCl}_2] - \Delta H_{\text{hyd}}^{\ominus}[\text{Mg}^{2+}]$ $\Delta H_{\text{hyd}}^{\ominus}[\text{Cl}^-] = -364 \text{ kJ mol}^{-1}$ Or $363.5 \text{ kJ mol}^{-1}$ [1 mark]	1
	c	Because the ionic radius of Mg^{2+} is smaller and there is less shielding than for Ba^{2+} . Or because the value of $\Delta H_{\text{hyd}}^{\ominus}$ is more exothermic for ions with the same charge but smaller ionic radii. Or $\Delta H_{\text{hyd}}^{\ominus}$ is more exothermic for Mg^{2+} than for Ba^{2+} because the charge density is greater for magnesium than for barium.	1

Part	Section	The answer	The mark
22	a(i)	HNO_3 or nitric acid	1
	a(ii)	Nitrobenzene	1
	b(i)	CH_3COCl or ethanoyl chloride	1
	b(ii)	Phenylethanone	1
	c	Bromoethane [1 mark] Bromoethane is more reactive than bromobenzene because the bond between carbon and bromine in bromoethane is weaker and easier to break. In bromobenzene, the bond is stronger due to the influence of the benzene ring, making it less reactive. [1 mark]	2
	d		1



Part	Section	The answer	The mark
23	a	$\text{HNO}_2 + \text{HCl}$ Or $\text{NaNO}_2 + \text{HCl}$ (1 mark for each)	2
	b	Phenylamine Or aniline	1
	c	 (1 mark) Sodium phenoxide (1 mark)	2
	d	The acidity of phenol is due to the delocalisation of the negative charge on the phenoxide ion into the π bonding electron system on the benzene ring. because it can lose a proton (H^+), to form phenoxide ion ($\text{C}_6\text{H}_5\text{O}^-$)	1

This is the end of the Marking Guide

