

This file was downloaded from the American Curriculum website



الملف Worksheet about Metal reactivity and properties and extraction and rusting

[Almanahj Website](#) → [American curriculum](#) → [9th Grade](#) → [Chemistry](#) → [Term 1](#) → [The file](#)

More files for 9th Grade , Subject Chemistry , Term 1

<a href="#">Chemistry exam preparation</a>	1
<a href="#">Solid state worksheet</a>	2
<a href="#">Chemistry Worksheet</a>	3
<a href="#">Worksheet about Hydrogen</a>	4
<a href="#">Mole to Grams and Grams to Moles Conversions Worksheet</a>	5
<a href="#">Worksheet about Chemistry</a>	6
<a href="#">Chemistry Worksheet</a>	7

Name: \_\_\_\_\_ ( ) Class: \_\_\_\_\_ Date: \_\_\_\_\_

**WORKSHEET**  
Chapter 14: Metals

**14.1 Metals and Alloys**


**Metals**

1. Metals are characterised by normally being shiny, strong solids with \_\_\_\_\_ melting points, which are \_\_\_\_\_ conductors of heat and \_\_\_\_\_.
2. Metals are \_\_\_\_\_ conductors of electricity because the \_\_\_\_\_ electrons in metallic atoms are not bound to any particular atom. The \_\_\_\_\_ ions of the metal are surrounded by a \_\_\_\_\_. Mobile electrons allow metals to conduct \_\_\_\_\_ and \_\_\_\_\_.
3. Metals have high \_\_\_\_\_, \_\_\_\_\_ and \_\_\_\_\_. This is because the atoms in metal are packed \_\_\_\_\_ in \_\_\_\_\_ and held together by \_\_\_\_\_. A large amount of \_\_\_\_\_ is required to break these bonds.
4. Metals are \_\_\_\_\_ and \_\_\_\_\_ because their atoms are of the same size and are closely packed in neat \_\_\_\_\_. When a \_\_\_\_\_ is applied, the layers \_\_\_\_\_ over one another.

**Alloys**

1. The properties of a particular metal can be improved if it is mixed with another element. We call these mixtures \_\_\_\_\_.
2. Compared to pure metals, \_\_\_\_\_ are \_\_\_\_\_ and \_\_\_\_\_, have a better \_\_\_\_\_, have \_\_\_\_\_ melting points and are \_\_\_\_\_ resistant to corrosion.

3. The atoms of a metal are arranged in regular \_\_\_\_\_. However if another element is added, this prevents the atoms from \_\_\_\_\_ one another. Thus, alloys tend to be \_\_\_\_\_ and \_\_\_\_\_ than pure metals.



Different sized atom of another element prevents atoms from \_\_\_\_\_ one another.

4. When zinc is added to copper the alloy formed is much \_\_\_\_\_ than pure copper. This alloy is called \_\_\_\_\_.
5. Another \_\_\_\_\_ of copper is \_\_\_\_\_ which consists of copper (90%) and tin (10%). The presence of \_\_\_\_\_ makes the copper \_\_\_\_\_ and less likely to corrode.

## 14.2 The Reactivity Series

1. In the reactivity series, metals are arranged from the \_\_\_\_\_ reactive to the \_\_\_\_\_ reactive.
2. Reaction of metals with cold water:

<b>Potassium</b>	Reacts _____. Highly _____. H <sub>2</sub> burns with a _____ flame. $2\text{K(s)} + 2\text{H}_2\text{O(l)} \rightarrow 2\text{_____ (aq)} + \text{H}_2\text{(g)}$
<b>Sodium</b>	Reacts _____. Highly _____. H <sub>2</sub> burns with a _____ flame. $2\text{Na(s)} + 2\text{H}_2\text{O(l)} \rightarrow 2\text{_____ (aq)} + \text{H}_2\text{(g)}$
<b>Calcium</b>	Reacts _____. $\text{Ca(s)} + 2\text{H}_2\text{O(l)} \rightarrow \text{_____}$
<b>Magnesium</b>	Reacts _____. $\text{Mg(s)} + 2\text{H}_2\text{O(l)} \rightarrow \text{_____}$
No reaction with zinc, iron, lead, copper and silver.	


3. Reaction of metals with steam:

<b>Potassium</b>	Reacts explosively. This reaction should not be carried out in the school laboratory.
<b>Sodium</b>	
<b>Calcium</b>	
<b>Magnesium</b>	Reacts _____ to produce a white glow. $\text{Mg(s)} + \text{H}_2\text{O(l)} \rightarrow \text{_____} + \text{H}_2\text{(g)}$
<b>Zinc</b>	Reacts _____. ZnO formed is yellow when hot, white when cold. $\text{Zn(s)} + \text{H}_2\text{O(l)} \rightarrow \text{_____}$
<b>Iron</b>	Reacts _____. $\text{Fe(s)} + 4\text{H}_2\text{O(l)} \rightarrow \text{Fe}_3\text{O}_4\text{(s)} + \text{H}_2\text{(g)}$
No reaction with lead, copper and silver.	

4. Reaction of metals with dilute hydrochloric acid:

<b>Potassium</b>	Reacts _____. $2\text{K(s)} + 2\text{HCl(aq)} \rightarrow 2\text{KCl(aq)} + \text{H}_2\text{(g)}$ $2\text{Na(s)} + 2\text{HCl(aq)} \rightarrow \text{_____}$
<b>Sodium</b>	
<b>Calcium</b>	Reacts _____. $\text{Ca(s)} + 2\text{HCl(aq)} \rightarrow \text{CaCl}_2\text{(aq)} + \text{_____}$
<b>Magnesium</b>	Reacts _____. $\text{Mg(s)} + 2\text{HCl(aq)} \rightarrow \text{_____} + \text{H}_2\text{(g)}$
<b>Zinc</b>	Reacts _____. $\text{Zn(s)} + 2\text{HCl(aq)} \rightarrow \text{_____}$
<b>Iron</b>	Reacts _____. $\text{_____}$
No reaction with lead, copper and silver.	

5. Metals at the top of the reactivity series, like \_\_\_\_\_ and \_\_\_\_\_ react violently with cold water. As you go down the series metals like \_\_\_\_\_ and \_\_\_\_\_ will only react with steam. Metals below lead, like \_\_\_\_\_, have no reaction at all with water or steam and will not displace \_\_\_\_\_ from water or steam.

	_____ <b>reactive</b>
	potassium
	calcium
	magnesium
	iron
	lead
	hydrogen
_____ <b>reactive</b>	

6. With dilute hydrochloric acid, the metals at the top of the series react very \_\_\_\_\_. As we go down the series, they react with decreasing ease. Metals below \_\_\_\_\_ in the series, like \_\_\_\_\_, will not displace hydrogen from dilute hydrochloric acid.

### 14.3 Using the Reactivity Series

- The reactivity series is related to the \_\_\_\_\_ of a metal to form \_\_\_\_\_ ions. Reactive metals tend to form \_\_\_\_\_ ions easily by \_\_\_\_\_ electrons and forming compounds. Unreactive metals prefer to remain uncombined as \_\_\_\_\_ of the element.
- One application of the reactivity series is the ability to predict the \_\_\_\_\_ power of a metal. A metal is capable of displacing another metal \_\_\_\_\_ in the reactivity series from its \_\_\_\_\_, or from its \_\_\_\_\_ solution.


3. Metal oxides that are \_\_\_\_\_ in the reactivity series are easier to reduce to the metal with carbon or hydrogen.
4. The \_\_\_\_\_ a metal is, the \_\_\_\_\_ it is to decompose its carbonate by heat. Carbonates of reactive metals are stable to heat. Carbonates of \_\_\_\_\_ metals decompose to form oxides of the metals and \_\_\_\_\_.

#### 14.4 Extracting Metals

1. Most metals are found in the ground combined with non-metals like \_\_\_\_\_, sulfur or carbon. These compounds are called \_\_\_\_\_.
2. Metals in their combined states, such as metal oxides, metal sulfides, metal chlorides and metal carbonates can be extracted by:
  - \_\_\_\_\_ with carbon, i.e. \_\_\_\_\_ the metal compound with carbon;
  - \_\_\_\_\_, which is using \_\_\_\_\_ to decompose the molten metal compound.
3. The stability of an ore of a metal higher up the reactivity series is \_\_\_\_\_ than one which is lower down. Therefore, it is much more \_\_\_\_\_ to extract sodium metal from its ore than it is to extract copper metal from its ore.
4. Every time a metal is extracted from its ore it involves \_\_\_\_\_ of the metal because the metal has to \_\_\_\_\_ electrons.
5. Metals which are \_\_\_\_\_ up in the reactivity series require strong reduction of their ores. This is usually achieved by \_\_\_\_\_. Middle order metals, such as \_\_\_\_\_, can be reduced by heating strongly with \_\_\_\_\_. Metals at the bottom of the reactivity series can be extracted from their ores simply by \_\_\_\_\_.
6. Some metals like \_\_\_\_\_ are so unreactive they occur in the ground as the metal itself.


## Extraction of Iron

1. Iron is extracted from its ores like \_\_\_\_\_ ( $\text{Fe}_2\text{O}_3$ ) in the blast furnace.
2. The furnace is charged at the top with three raw materials — \_\_\_\_\_, \_\_\_\_\_ and \_\_\_\_\_.
- 3.




## Chemical Reactions in the Blast Furnace

1. Production of carbon dioxide  
Coke burns in hot air. Carbon dioxide is produced. Heat is given off as the reaction is highly \_\_\_\_\_ and the temperature in this part of the furnace rises to  $1900^\circ\text{C}$ .




2. Production of carbon monoxide  
Carbon dioxide reacts with hot \_\_\_\_\_ to produce \_\_\_\_\_.



3. Reduction of iron ore

Carbon monoxide gas is a powerful \_\_\_\_\_ and it reduces the iron ore at the top of the furnace to molten iron. The molten iron falls to the bottom of the furnace where it is tapped off.



4. Removal of impurities

The \_\_\_\_\_ present in the furnace helps to remove impurities. The main impurity in the iron is sand (\_\_\_\_\_). Inside the furnace, the \_\_\_\_\_ (calcium carbonate) decomposes to \_\_\_\_\_ (calcium oxide) and \_\_\_\_\_.



Calcium oxide reacts with \_\_\_\_\_ from sand in iron ore to form calcium silicate, also known as \_\_\_\_\_. Slag floats on top of \_\_\_\_\_ and is used in \_\_\_\_\_.



Carbon dioxide escapes as \_\_\_\_\_ from the top of the furnace together with \_\_\_\_\_ and \_\_\_\_\_.

### 14.5 Uses of Steel


1. The iron produced in the blast furnace is \_\_\_\_\_ and contains impurities of mainly \_\_\_\_\_ and other non-metals. It is called \_\_\_\_\_ or pig iron. Such iron is very brittle and has limited uses. Most of this iron is converted into \_\_\_\_\_.
2. Steel is produced in a \_\_\_\_\_. Here under high pressure \_\_\_\_\_ is blown into the molten cast iron. This oxidises the \_\_\_\_\_ like carbon and sulfur to gases which then escape. Other impurities like phosphorus and silicon are converted into \_\_\_\_\_ which can be neutralised by the addition of a base such as \_\_\_\_\_.




3. When all the impurities have been \_\_\_\_\_ we are left with a very pure form of iron called \_\_\_\_\_.
4. Calculated amounts of \_\_\_\_\_ are then added to produce the various steels. \_\_\_\_\_ is low carbon steel and contains about \_\_\_\_\_ carbon. It is \_\_\_\_\_ and therefore easy to shape. It is ideal for making \_\_\_\_\_ and machinery.
5. The more carbon that is added, the \_\_\_\_\_ the steel becomes. This is because the carbon impurity strengthens the iron lattice, making it more difficult for the atoms to \_\_\_\_\_ one another. High carbon steel contains between 0.5 to 1.5% carbon and is \_\_\_\_\_ but \_\_\_\_\_. It is used for making \_\_\_\_\_ and \_\_\_\_\_ tools.
6. Other alloys of steel are also made by the addition of the correct amount of the appropriate metal. Stainless steel contains iron, carbon, \_\_\_\_\_ and \_\_\_\_\_. It is attractive and very resistant to \_\_\_\_\_. It is therefore useful in making \_\_\_\_\_, \_\_\_\_\_ and equipment in chemical plants.

#### 14.6 Rusting

1. The essential conditions for the rusting ( \_\_\_\_\_ ) of iron are the presence of \_\_\_\_\_ and \_\_\_\_\_. This results in the flaky brown coating, commonly called rust but chemically known as \_\_\_\_\_ ( $\text{Fe}_2\text{O}_3 \cdot x\text{H}_2\text{O}$ ). The presence of \_\_\_\_\_ and \_\_\_\_\_ substances speed up rusting.



2. In which test tube will the nails be most heavily rusted after one week?



	Rust		Rust		Rust		Rust
	No rust		No rust		No rust		No rust

The nails in test tube \_\_\_ will be most heavily rusted.

- Prevention of rusting can be achieved by placing a barrier around the metal. Examples are \_\_\_\_\_, greasing, covering with \_\_\_\_\_ and \_\_\_\_\_.
- An alternative way of preventing rusting is by \_\_\_\_\_. A more reactive metal (like magnesium or \_\_\_\_\_) is attached to the iron object. The more reactive metal \_\_\_\_\_ in preference to the iron.
- Sacrificial protection is useful with underground steel or iron objects like \_\_\_\_\_ or \_\_\_\_\_ in petrol stations. These are difficult to paint and grease, but it is easy to attach a piece of \_\_\_\_\_ or \_\_\_\_\_ to such objects.
- Rusting can also be prevented by using rust-resistant alloys such as \_\_\_\_\_.